



# Quantification of chlorogenic acid in *Pluchea indica* L. stem ethanolic extracts and its antioxidant activity

NI PUTU ERMI HIKMAWANTI\*, AGUSTIN YUMITA, JIHAN ESA SIREGAR

Department of Pharmaceutical Biology, Universitas Muhammadiyah Prof. dr Hamka, East Jakarta, Indonesia

## ARTICLE INFO

Received 17 March 2023  
Accepted 28 August 2023

### Keywords:

Beluntas,  
chlorogenic acid,  
DPPH,  
ultrasonic-assisted extraction,  
sequential extraction.

## ABSTRACT

Chlorogenic acid (CA) is an important phenolic acid antioxidant. It is found in *Pluchea indica* L. (Asteraceae). However, it has only been extensively studied in the leaves, while studies on the stems have not been reported. This study aimed to identify and measure the levels of CA in the stem extract of *P. indica*. The extract was also determined for its antioxidant activities. In the course of the work, *P. indica* stems powder was extracted using the ultrasonic-assisted extraction (UAE) technique employing 50%-ethanol as solvent directly and sequentially. The extract was then measured for total phenolic content (TPC) and CA content using RP-HPLC. Meanwhile, antioxidant activities were determined by the DPPH, ABTS, and reducing power (RP) methods. TPC in the sequential and the direct of *P. indica* stems ethanol extracts were  $1.4694 \pm 0.0228$  and  $1.9314 \pm 0.0318$  mgGAE/g DW, respectively. We found that the CA content of 50%-ethanol extract of *P. indica* stems from sequential extraction ( $0.2045 \pm 0.0128\%$ , w/w) was higher than 50%-ethanol extract from direct extraction ( $0.1984 \pm 0.0113\%$ , w/w). The two extracts demonstrated good antioxidant capacity, while the ethyl acetate and *n*-hexane extracts did not. Identifying of other antioxidants phenolics using other extracting methods still needs further study.

## INTRODUCTION

Chlorogenic acid (CA) is an important phenolic derivative of caffeoylquinic acid (CQA). CA (or 5-CQA) consists of a quinic acid core which is acylated with one moiety of caffeic acid at the C-5 (Figure 1) [1]. CA is applicable as an antioxidant, anti-inflammatory, antidiabetic, anti-obesity, anti-hypertensive, antimicrobial, etc.[1]. The caffeic acid group in CA plays a critical role in its activity as an antioxidant. According to Kritsi *et al.* [2], CA has antioxidant activity through the interaction of hydrogen bonds on the receptor binding sites of NADPH oxidase, cytochrome P450, and myeloperoxidase [2]. CA can be extracted from plant materials using ultrasound-assisted extraction (UAE) [3]. CA can be analyzed qualitatively or quantitatively using High-performance liquid chromatography (HPLC) [4] with a C18 column with a UV or DAD detector [4].

CA is found in many plants [3]. One source of CA from plants is *Pluchea indica* L. (Asteraceae), especially the leaves [5-8]. The *P. indica* leaves were more explored than other plant parts. *P. indica* stems also contained a relatively high total phenolic content in 50%-ethanol extract compared to flowers and roots [9]. The importance of CA

compounds as a source of antioxidants that can reduce free radicals needs to be explored. Indeed, the quantification of CA in 50%-ethanol extract of *P. indica* stems has never been reported. In the present study, quantification of the compound in *P. indica* stems was carried out as an ethanol extract (obtained directly and sequentially), using RP-HPLC analysis. Subsequently, the antioxidant activities of the extracts were studied.




Figure 1. Chlorogenic acid (5-CQA)

## MATERIALS AND METHODS

### Chemicals

CA and gallic acid as reference were purchased from MarkHerb, Institut Teknologi Bandung (ITB), Bandung, Indonesia. Ethanol, ethyl acetate, and *n*-hexane (PT. Brataco, Indonesia) were used as extraction solvents. Methanol, acetic acid, and water for HPLC grade (Merck, Germany) were used for HPLC analysis.

\* Corresponding author

e-mail: [ermy0907@uhamka.ac.id](mailto:ermy0907@uhamka.ac.id)

## Plant material

*P. indica* stems were collected on November 2021 from the Biopharmaceutical Cultivation Conservation Unit, Tropical Biopharmaca Research Center (TropBRC), Institut Pertanian Bogor (IPB), Bogor, Indonesia, with voucher number: BMK0188092016. The stems were cleaned and air-dried. The dried stems were then powdered and stored in a dry container protected from light.

## Extraction

Extraction is divided into two ways, namely direct extraction and sequential extraction. In our work, each five g of stem-dried powder was extracted with 50 mL of solvent. In summary, the extraction procedure is presented in Figure 2. The extraction process was carried out using the UAE technique following the protocol in Kongkiatpaiboon *et al.* [6], and was performed in an ultrasonic bath (Branson 5510) 40 kHz at 40°C for 15 min, with the filtrate being separated from the residue by filtration. The residue was then re-extracted using new solvent 3 times. Each filtrate was subsequently collected and made up to 50 mL (here-to-after, referred to as 'liquid extract'). Extraction of each sample was carried out for 5 replications.




Figure 2. Extraction procedures of *P. indica* dried stem material

## Polyphenol Screening

Screening of phenolic compounds was carried out qualitatively using a 5% FeCl<sub>3</sub> reagent. The positive result of the reaction is indicated by the formation of a blue-black colour [10].

## Total Phenolic Content Assay

Colorimetry was used to determine total phenolic content following the Hikmawanti *et al.* [11] procedure with slight modifications. Gallic acid was used as a standard. For calibration, the gallic acid solution was prepared with variation concentrations:  $1.1 \times 10^{-2}$ ,  $1.7 \times 10^{-2}$ ,  $2.3 \times 10^{-2}$ ,  $2.9 \times 10^{-2}$ , and  $3.5 \times 10^{-2}$  mg/mL. Each extract was diluted 20 times using ethanol. Both extract and standard (0.3 mL) were mixed with 1.5 mL of the Folin-ciocalteu (1:10 in water) and left for 3 min. Afterwards, 1.2 mL of sodium carbonate solution (7.5%) was added to the mixture, homogenized, and then incubated at room temperature for 110 min. The absorbance was measured at 765.10 nm with a spectrophotometer UV-Vis (UV-1900i Series, Shimadzu, Kyoto, Japan).

The total phenolic content was expressed as gallic acid equivalent (mg GAE/g of the dried sample). The test was carried out in triplet (mean  $\pm$  SD).

## HPLC condition for analysis of CA

CA analysis followed the protocol of Kongkiatpaiboon *et al.* [6] with slight modification, and was undertaken at The Advanced Characterization Laboratories Cibinong-Integrated Laboratory of Bioproduct, National Research and Innovation Agency (BRIN), Indonesia. CA (as an external standard) was prepared at a concentration of 10 mg/mL in ethanol. The standard solution was then diluted to 1.0, 2.0, 6.0, and 7.5 mg/mL. Each solution was filtered using a 0.45  $\mu$ m filter and injected into the HPLC system. The injection volume of the sample was 5  $\mu$ L. The HPLC system (Shimadzu Prominence-i LC-2030C 3D Plus), which includes a quaternary pump, a degasser and an autosampler, is combined with a photodiode array (PDA) detector in the HPLC apparatus. HPLC separation was performed using a Shim Pack GIST C18 column (150 mmL.  $\times$  4.6 mm I.D., 4  $\mu$ m) as the stationary phase at 25°C. Acetic acid 0.5% in water (A) and methanol (B) was used as the mobile phase via gradient elution (10% B to 50% B in A for 40 min and 100% B for 10 min), as stated in Kongkiatpaiboon *et al.* [6]. The flow rate was 1.0 mL/min. Separation was monitored at 326 nm.

## DPPH assay

A 0.1 mM 2,2-Diphenyl-1-picrylhydrazyl (DPPH) solution was prepared in methanol. Quercetin ( $1 \times 10^2$  mg/mL) in methanol was used as a reference. Initially, each sample of both extract and quercetin (1 mL) was separately reacted with 0.1 mM DPPH (4 mL). After incubation for 30 min at room temperature in the dark, the reaction of the mixture was measured at 516 nm against a methanol blank, using a UV-Vis spectrophotometer (UV-1900 Series, Shimadzu, Kyoto, Japan) [12].

## ABTS assay

Antioxidant activity assay against 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS) radicals was carried out by microtiter assay. Accordingly, 180  $\mu$ L of ABTS radical (ABTS<sup>•+</sup>), prepared 12-16 h beforehand, was reacted with 20  $\mu$ L of the extract test solution in a 96-well flat-bottom microplate. After 5 min of incubation, the absorbance was measured at 750 nm, using an iMark microplate reader (BioRad, Shanghai, China) [13]. Trolox was employed as a reference (0.1-0.8 mmol/L). The results are expressed as  $\mu$ mol Trolox equivalent per g dry weight sample ( $\mu$ mol TE/g DW).


## RP assay

The ability of antioxidants to reduce iron was determined by reducing power (RP) assay following the protocol of Khatua *et al.* [13]. The tests were carried out in 96-well flat-bottom microplates. A total of 10  $\mu$ L of the extract test solution was reacted with 25  $\mu$ L of 0.2 M phosphate buffer (pH 6.6), and 25  $\mu$ L of potassium ferricyanide (1%, w/v). After 20 min incubated (at room temperature), 25  $\mu$ L of Trichloroacetic Acid (TCA) 10% (w/v) was added.

Subsequently, 5  $\mu$ L of distilled water and 8.5  $\mu$ L of Iron (III) chloride was added in. The mixture was shaken for 10 s at medium speed, then incubated at room temperature for 15 min. The absorbance was measured at 750 nm using a iMark microplate reader (BioRad, Shanghai, China) [13]. Trolox was employed as a reference (0.1-0.8 mmol/L). The results are expressed as  $\mu$ mol Trolox equivalent per g dry weight sample ( $\mu$ mol TE/g DW).

## RESULTS

The 50%-ethanol extracts of *P. indica* stems, both obtained from direct and sequential extraction, were detected to contain phenolics based on the test with a 5%  $\text{FeCl}_3$  reagent (Figure 3). The two extracts were then analysed for the total phenolics content via a UV-Visible spectrophotometer. Total phenolics content in the 50%-ethanol extract of *P. indica* stems from sequential and direct extraction were  $1.4694 \pm 0.0228$  and  $1.9314 \pm 0.0318$  mgGAE/g dried powder, respectively.




A - 50%-ethanol extract (direct); B - 50%-ethanol extract (sequential); C - ethyl acetate extract (sequential); D - *n*-hexane extract (sequential)

**Figure 3.** The results of phenolic compounds screening in *P. indica* stem extracts with 5%  $\text{FeCl}_3$  reagent

Analysis of CA in ethanol extracts of *P. indica* stems was performed using RP-HPLC at 326 nm as the maximum wavelength [6]. Figure 4 shows the HPLC chromatograms of the standard CA (A, retention time = 23.042 min), the 50%-ethanol extract (sequential) (B), and the 50%-ethanol extract (direct) (C) of *P. indica* stems. The chromatogram showed the presence of CA in the 50%-ethanol extract (sequential) and the 50%-ethanol extract (direct) of *P. indica* stems at retention times of 22.702 dan 22.946 min, respectively. The equation of the line obtained from the CA calibration curve is  $y = 1628x - 3E + 06$  ( $R^2 = 0.9991$ ). The quantification of CA in the 50%-ethanol extract (sequential) and the 50%-ethanol extract (direct) of *P. indica* stems were  $0.2045 \pm 0.0128$  and  $0.1984 \pm 0.0113\%$  (w/w), respectively.

The antioxidant activity of the stem extract of *P. indica* was tested by way of application of the DPPH, ABTS and RP methods. Table 1 shows the summary of the antioxidant activity of the extracts. The two ethanol extracts (direct and



A - chlorogenic acid; B - 50%-ethanol extract (sequential); C - 50%-ethanol extract (direct)

**Figure 4.** RP-HPLC Chromatogram of chlorogenic acid (CA) as a standard (at 2.0 mg/mL) and ethanolic extracts of *P. indica* stem

sequential) provide an overview of the activity of reducing radicals (both DPPH $\cdot$  and ABTS $^{+}$ ) and metal ions (through iron reduction). The ethyl acetate and *n*-hexane extracts did not provide an overview of antioxidant capacity.

**Table 1.** Antioxidant activity from *P. indica* stem extracts

Samples	Antioxidant activities		
	DPPH (mg QE/100 g DW)	ABTS ( $\mu$ mol TE/g DW)	RP ( $\mu$ mol TE/g DW)
50%-ethanol extract (direct)	$189.210 \pm 0.01$	$17.624 \pm 0.345$	$40.069 \pm 0.214$
50%-ethanol extract (sequential)	$143.198 \pm 0.02$	$34.837 \pm 0.126$	$42.854 \pm 0.909$
Ethyl acetate extract (sequential)	$1.209 \pm 0.05$	nd	nd
<i>n</i> -hexane extract (sequential)	$0.597 \pm 0.03$	nd	nd

Note: DW - dry weight; nd - not detected; QE - quercetin equivalent; TE - Trolox equivalent

## DISCUSSION

CA is a compound biosynthesized through the shikimate pathway in plants from various families, including Asteraceae [14]. It is accumulated in leaves, stems, roots, fruits and other organs at different levels [15]. Extraction is the initial step in separating metabolites from the plant matrix for identification [16]. To measure the quantity of CA in *P. indica* extract, the HPLC technique might be used for routine analysis [6]. This method is proven to be fast, precise and accurate. In previous study, it was reported that CA, as one of the phenolics of *P. indica* leaves, can be efficiently extracted using 50%-ethanol solvent with UAE [6]. In the present study, phenolic extraction was performed using 50%-ethanol solvent, directly and sequentially, by way of use of the UAE technique. CA was detected and quantified

in 50%-ethanol extract (both obtained from direct and sequential extraction). Compared with the levels of CA in *P. indica* leaves in the study of Kongkiatpaiboon *et al.* [6], the content of this compound in the ethanol extract of *P. indica* stems was lower. The levels of CA in *P. indica* leaves were in the range of 2.67% (w/w) [6], while that in stem extracts (in the present study) were in the range of 0.19-0.2% (w/w). Direct extraction allows metabolites with a polarity corresponding to 50%-ethanol to be extracted in it (crude), thus the concentration of the target compound in this extract is lower. Through sequential extraction, it is expected that other compounds extracted sequentially in *n*-hexane and ethyl acetate are not found in the 50%-ethanol extract. With the elimination of these compounds (such as components of fat, chlorophyll, etc.), it is hoped that phenolics (especially CA) can be effectively extracted in 50%-ethanol solvent [17]. In general, CA is soluble in low concentrations of alcohol or alcohol-water mixtures. This compound is insoluble in non-polar solvents, such as benzene, chloroform, or ether. The high solubility in an alcohol-water mixture is related to the large number of free hydroxyl groups [18].

In this study, the high phenolic content in the extract was linear with its antiradical DPPH activity. The 50%-ethanol extract (direct) had high phenolic content and the DPPH radical scavenging activity was better than that with other extracts. The extract had, however, a lower CA content than the 50%-ethanol extract (sequential). Thus, it is suspected that CA is not the only phenolic in *P. indica* stems that plays a role in scavenging DPPH radicals. The presence of other phenolic compounds, or other natural compounds that are synergistic in the 50%-ethanol extract, is suspected of playing a role in providing better antiradical activity than the 50%-ethanol extract (sequential). In the ABTS and RP tests, 50%-ethanol extract (sequential) gave better antioxidant activity.

There are several chemical approaches for measuring plant antioxidant capacity. The antioxidant chemical-based assay's mode of action is that of electron transfer, hydrogen transfer and metal reduction [19]. The DPPH and ABTS methods are included in reactions based on electron and hydrogen transfer. The DPPH technique is an *in vitro* approach that is simple, inexpensive and frequently used to assess the antioxidant activity of natural materials [20]. This method describes the antiradical activity of the sample more than as an antioxidant activity. Antiradical activity refers to a compound's capacity to react with a free radical, whereas antioxidant activity refers to a compound's ability to block the oxidation process [21]. Meanwhile, ABTS is capable of measuring the capacity of both lipophilic and hydrophilic antioxidants. ABTS radicals are more reactive than DPPH radicals. The chemical interaction between antioxidants and ABTS radicals is relatively quick [22]. Another method, RP, is based on an increase in absorbance from the reaction between the antioxidant and a mixture of potassium ferricyanide, trichloro acetic acid and ferric chloride reagents, which demonstrates the strength of antioxidants in reducing metals [20]. However, measuring antioxidant activity *in vitro* has limitations, since DPPH and ABTS are not naturally occurring oxidants in the body. The radical quenching

reactions that occur in these assays are often quick, unlike in the occurrence of a slower reaction in the body. In addition, whereas antioxidants have significant antioxidant effects in chemical tests, they can function as pro-oxidants in dietary and biological systems [19].

The variety of phenolic types and the complexity of the composition of these chemicals in plants make the separation of antioxidants phenolic inefficient and expensive [17]. Other compounds that can also act as antioxidants are phytoosterols [23], flavonoids, minerals, organosulfur, vitamins, carotenoids, etc. [24]. Moreover, the crude often provides more significant antioxidant benefits than the individual forms. In addition, the measurement of antioxidant activity with specific methods is usually only carried out under adjusted conditions. Determining the overall antioxidant capacity of the sample should, therefore, be also done using additional techniques, exceptionally those capable of evaluating the efficiency of antioxidants against specific ROS/RNS [17]. Studying the types of other phenolic compounds in *P. indica* stems is still possible. Thus, the use of *P. indica* stems as a source of antioxidants can continue to be developed, for example, in the therapy of metabolic syndrome, as a preservative, and as natural food supplements.

## CONCLUSIONS


Based on this research, the total phenolic content in the 50%-ethanol extract of *P. indica* stems from direct extraction is more than the 50%-ethanol extract obtained from sequential extraction. Meanwhile, the 50%-ethanol extract from sequential extraction contains more CA than the 50%-ethanol extract obtained from direct extraction. The extracts also provided better antioxidant activity. It is important, therefore, to search for phenolic substances besides CA that function as antioxidants in *P. indica* stems.


## ACKNOWLEDGEMENT


The authors wish to thank the Research and Development Institute or Lembaga Penelitian dan Pengembangan (Lemlitbang), Universitas Muhammadiyah Prof. DR. HAMKA for funding this research and publication with grant number: 713/F.03.07/2021. The authors also thank the facilities, scientific and technical support from Advanced Characterization Laboratories Cibinong-Integrated Laboratory of Bioproduct, National Research and Innovation Agency through E-Layanan Sains (ELSA), Badan Riset dan Inovasi Nasional (BRIN), Indonesia.

## ORCID iDs

Ni Putu Ermi Hikmawanti

 <https://orcid.org/0000-0001-5194-1431>

Agustin Yumita  <https://orcid.org/0000-0003-2037-2648>

Jihan Esa Siregar  <https://orcid.org/0000-0003-0220-9484>

## REFERENCES

1. Naveed M, Hejazi V, Abbas M, Kamboh AA, Khan GJ, Shumzaid M, et al. Chlorogenic acid (CGA): A pharmacological review and call for further research. *Biomed Pharmacother.* 2018;97:67-74.
2. Kritsi E, Tsiaka T, Ioannou AG, Mantanika V, Strati IF, Panderi I, et al. *In Vitro* and *in silico* studies to assess edible flowers' antioxidant activities. *Appl Sci.* 2022;12(14):7331.

3. Lu H, Tian Z, Cui Y, Liu Z, Ma X. Chlorogenic acid: A comprehensive review of the dietary sources, processing effects, bioavailability, beneficial properties, mechanisms of action, and future directions. *Compr Rev Food Sci Food Saf*. 2020;1-29.
4. Wianowska D, Gil M. Recent advances in extraction and analysis procedures of natural chlorogenic acids. *Phytochem Rev*. 2019;18(1): 273-302.
5. Arsiningtyas IS, Gunawan-Puteri MDPT, Kato E, Kawabata J. Identification of  $\alpha$ -glucosidase inhibitors from the leaves of *Pluchea indica* (L.) Less., a traditional Indonesian herb: promotion of natural product use. *Nat Prod Res*. 2014;28(17):1350-3.
6. Kongkiatpaiboon S, Chewchinda S, Vongsak B. Optimization of extraction method and HPLC analysis of six caffeoylquinic acids in *Pluchea indica* leaves from different provenances in Thailand. *Rev Bras Farmacogn*. 2018;28(2):145-50.
7. Vongsak B, Kongkiatpaiboon S, Jaisamut S, Konsap K. Comparison of active constituents, antioxidant capacity, and  $\alpha$ -glucosidase inhibition in *Pluchea indica* leaf extracts at different maturity stages. Vol. 25, *Food Bioscience*. 2018. p. 68-73.
8. Chewchinda S, Vongsak B. Simultaneous HPTLC quantification of three caffeoylquinic acids in *Pluchea indica* leaves and their commercial products in Thailand. *Rev Bras Farmacogn*. 2019;29: 177-81.
9. Normala H, Suhaimi MI. Quantification of total phenolics in different parts of *Pluchea indica* (Less) ethanolic and water extracts. *Pertanika J Sci Technol*. 2011;19(1):19-24.
10. Shaikh JR, Patil M. Qualitative tests for preliminary phytochemical screening: An overview. *Int J Chem Stud*. 2020;8(2):603-8.
11. Hikmawanti NPE, Hanani E, Sapitri Y, Ningrum W. Total phenolic content and antioxidant activity of different extracts of *Cordia sebestena* L. Leaves. *Pharmacogn J*. 2020;12(6):1311-6.
12. Molyneux P. The Use of the stable free radical diphenylpicrylhydrazyl (DPPH) for estimating antioxidant activity. *Songklanakarin J Sci Technol*. 2004;26(2):211-9.
13. Khatua S, Ghosh S, Acharya K. Simplified methods for microtiter based analysis of in vitro antioxidant activity. *Asian J Pharm*. 2017; 11(2):S327-35.
14. Dewick PM. *Medicinal natural products: A Biosynthetic Approach*. John Wiley & Sons, Ltd; 2002:1-539.
15. Liu W, Li J, Zhang X, Zu Y, Yang Y, Liu W, et al. Current advances in naturally occurring caffeoylquinic acids: Structure, bioactivity, and synthesis. *J Agric Food Chem*. 2020;68(39):10489-516.
16. Zhang QW, Lin LG, Ye WC. Techniques for extraction and isolation of natural products: A comprehensive review. *Chin Med*. 2018;13(20).
17. Dai J, Mumper RJ. Plant phenolics: Extraction, analysis and their antioxidant and anticancer properties. *Molecules*. 2010;15:7313-52.
18. Upadhyay R, Mohan Rao LJ. An outlook on chlorogenic acids-occurrence, chemistry, technology, and biological activities. *Crit Rev Food Sci Nutr*. 2013;53(9):968-84.
19. Huang D, Wong IC. Antioxidant evaluation and antioxidant activity mechanisms. In: Logan A, Nienaber U, Pan X (ed). *Lipid oxidation: Challenges in food systems*. Academic Press and AOCS Press; 2013:323-43.
20. Alam MN, Bristi NJNJ, Rafiquzzaman M. Review on *in vivo* and *in vitro* methods evaluation of antioxidant activity. *Saudi Pharm J*. 2013;21(2):143-52.
21. Tirezitis G, Bartosz G. Determination of antiradical and antioxidant activity: basic principles and new insights. *Acta Biochim Pol*. 2010; 57(1):139-42.
22. Gulcin I. Antioxidants and antioxidant methods: an updated overview. *Arch Toxicol*. 2020;94(3):651-715.
23. Wang T, Hicks KB, Moreau R. Antioxidant activity of phytosterols, oryzanol, and other phytosterol conjugates. *JAACS*. 2002;79(12): 1201-6.
24. Munteanu IG, Apetrei C. Analytical methods used in determining antioxidant activity: A review. *Int J Mol Sci*. 2021;22(7).