


Current Issues in Pharmacy and Medical Sciences

Formerly ANNALES UNIVERSITATIS MARIAE CURIE-SKŁODOWSKA, SECTIO DDD, PHARMACIA

journal homepage: <http://www.curipms.umlub.pl/>



Bioactivity-guided isolation of alkamides from a cytotoxic fraction of the ethyl acetate extract of *Anacyclus pyrethrum* (L.) DC. roots

SOUAD HAMIMED^{1,2,5}, NADJI BOULEBDA³, HOCINE LAOUER⁴, ABDELMALIK BELKHIRI^{1,2*}

¹ Laboratoire de Pharmacognosie, Faculté de Médecine, Université Salah Boubnider, Constantine 3, Algérie

² Laboratoire de Pharmacologie and Toxicologie, INSV – Khroub, Université Mentouri Constantine 1, Algérie

³ Laboratoire de Pharmacologie, ISAV – Université Mohamed-Cherif Messaadia – Souk Ahras, Algérie

⁴ Laboratoire de Valorisation des Ressources Naturelles Biologiques. Université Ferhat Abbes Sétif, Algérie

⁵ Université Larbi Ben M'hidi O.E.B, Algérie

ARTICLE INFO

Received 08 February 2018

Accepted 10 May 2018

Keywords:

Artemia salina,
DL50,
fractionation,
alkamides,
pellitory.

ABSTRACT

Introduction. The alcohol extract of Pellitory (*Anacyclus pyrethrum*) roots has been previously shown to exert anticancer activities on the Human Colorectal Cancer Cell Line (HCT) by targeting apoptosis, metastasis and cell cycle arrest. However, the nature of the cytotoxic molecules associated with this activity remains unexplored.

Aims. This study aims to reinvestigate Pellitory root extract as regard to its cytotoxic activity and to proceed to a bioguided fractionation to explore its active fraction and to give new insight in their phytochemical constituents.

Methods. Powdered roots were subjected to repeated extraction with Petroleum ether (Pe), Chloroform (Ch), Ethyl acetate (Ea) and Methanol (Me). Pellitory extracts were then screened for cytotoxic activity using the Brine Shrimp Lethality (BSL) bioassay.

Results. Ea extract exhibited a marked cytotoxic activity, with LC50 of 249.26 µg/mL in the BSL bioassay. The remaining extracts (Pe,Ch,Me) treated groups exhibited no or low mortality in the range of tested concentrations (1-1000 µg/mL). BSL assay-guided chromatographic fractionation of Ea active Extract revealed a highly cytotoxic fraction (F11) with LC50 of 42.5 µg/mL. Multistep purifications of the active F11 fraction afforded four alkamides, namely N-isobutyldeca-2,4-dienamide or Pellitorine (I), N-propyldodeca-2,8-dienamide (II), N-isobutyltetradeca-2,4-dienamide (III) and N-propylnona-2,5-dienamide (IV).

Conclusions. This study suggests that cytotoxic activity is localized mainly in the ethyl acetate extract (Ea) of pellitory roots. BSL assay fractionation of this active extract leads to the isolation of four alkamides, including pellitorine (I). While this isobutyl alkamide has previously shown strong cytotoxic activities against human cancer cell lines, the other compounds (II to IV) were not previously reported as cytotoxic. Subsequently, the isolated alkamides will be considered in future study as candidates for in depth in-vitro evaluation of their cytotoxicity against cancer and normal cell lines. Finally, through this study, BSL assay demonstrate again its usefulness as bench-top assay in exploring plant extracts for cytotoxic compounds.

Abbreviations

BSL (Brine Shrimp Lethality), Ch (Chloroform), DMSO (Dimethyl sulfoxide), Ea (Ethyl acetate), HCT (Human Colorectal Cancer Cell Line), MCT-7 (Brest cancer strain),

Me (Methanol), HL60 (Human promyelocytic leukemia cell line), Pe (Petroleum ether).

INTRODUCTION

Pellitory (*Anacyclus pyrethrum* (L.) DC., syn.: *Anacyclus officinarum* Hayne), is a perennial procumbent herb which

* Corresponding author

e-mail: abdelmalik.belkhiri@univ-constantine3.dz

belongs to the Asteraceae family. The species is widely distributed in North Africa, and elsewhere in the Mediterranean region, in India and in Arabian countries [1]. The root of pellitory is well-known as a medicinal drug. It is used in the ayurvedic and unani systems of medicine for its medicinal properties, such as anti-rheumatic, odontalgia, antibacterial, antiviral, carminative, anti-catarrah, digestive, emmenagogue, febrifuge, vermifuge, sialagogue and anti-cancer [2-8]. In the North Africa region, this herbal drug is known as “*Guentess*”, and is indicated among other therapeutic applications, in respiratory infections and in the treatment of hepatic disorders [9-11].

Scientific studies has revealed that extracts from the root exhibit antibacterial, antinociceptive, anti-inflammatory, immunostimulating and antioxidant activities [7,12,13]. Moreover, the phytochemical screening of *Anacyclus pyrethrum* has led to the identification of various secondary metabolites such as alkaloids, tannins, flavonoids, coumarins and lignanes [12]. This species also contains saponins, sesamin, inulin, gum and traces of essential oil [13]. The most important chemical markers present in its root are N-alkamides with pellitorine and polysaccharides [14-16]. Furthermore, LC-MS N-alkamides profiling of an ethanolic *Anacyclus pyrethrum* root extract reported thirteen N-alkamides, including N-isobutyldeca-2,4-dienamide (Pellitorine) and N-isobutyltetradeca-2,4-dienamide [14].

Recent studies showed that *Anacyclus pyrethrum* extract have cytotoxic activity, and can successfully induce apoptosis in human colorectal cancer lines HCT cells [17,18]. Although, these experimental results give relatively some credence to the reported ethnobotanical use of the plant as anticancer [8], the identity of the active compounds associated with this effect is still unknown so far.

On the basis of this information, we decided to investigate *Anacyclus pyrethrum* roots extracts using brine shrimp lethality (BSL) assay as a predictive test for cytotoxicity [19]. This bioassay will be used to monitor the fractionation steps of the active extract and give new insight in its bioactive fractions and their phytochemical constituents. The isolated compounds can subsequently be tested in future studies using specific antitumor assays of interest.

MATERIALS AND METHODS

Solvents and Reagents

All substances were purchased from Sigma-Aldrich Chemical Co (Strasbourg, France) unless otherwise stated.

Plant material

Anacyclus pyrethrum (L) DC. (Asteraceae) roots were collected nearby Constantine, Algeria. The plant was identified by a Taxonomist (Pr.H. Laouer, Setif University) and voucher specimen (Ref. AN-00301R) was deposited for

future reference in the herbarium at the laboratory of botany, University SB Constantine 3, Algeria.

Extraction method

Pellitory extracts were obtained from the air-dried roots (250 g finely ground), by successive maceration (2×1.5 L) with petroleum ether (Pe), chloroform (Ch), ethyl acetate (Ea) and methanol (Me). The extractive solutions were evaporated to dryness under vacuum to obtain 22.8 g (Pe), 18.0 g (Ch), 29.8 g (Ea) and 56.3 g (Me) extracts. Table 1 shows the corresponding yields of the crude extracts, expressed as percentage (w/w) dry powder.

Table 1. Cytotoxicity profile of pellitory extracts

Treatment	Symb	Yield# (%)	Mortality, expressed as percentage Δ						LC 50 (μg/mL)	Toxicity profile ¥	
			Concentration (μg/mL)								
			1	10	50	100	250	500			1000
Petroleum ether	(Pe)	9.12	-	-	-	-	-	10 ±0.5*	20.0 ±0.5**	2044.65	n/T
Chloroform	(Ch)	7,60	-	-	-	-	-	-	10.0 ±0.0	1391.69	n/T
Ethyl acetate	(Ae)	11.92	-	-	-	10.0 ±0.0	25.0 ±0.6***	100.0 ±0.0	100.0 ±0.0	249.26	T
Methanol	(Me)	22.52	-	-	-	-	-	-	47,5 ±1.0***	1000.0	n/T

#: yield (Percentage w/w, dry matter)

Δ: Values of mortality after 24h, expressed as mean +/- SD (n=4)

(*p<0.05, **p<0.01, ***p<0.001): value vs negative control (1% DMSO)

-: no mortality of nauplii was recorded

¥: LC 50 ≥1000 μg/ml is considered nontoxic (n/T). LC50<1000 μg/ml is toxic (T)

Phytochemical screening

Chemical screening of crude extracts, fractions and sub-fractions were carried out using well-established staining and precipitation reactions for major groups of bioactive natural products [20].

Thin layer chromatography (TLC) control of fractions and isolated compounds was carried out on precoated silica gel 60F₂₅₄ aluminium plates (5×10 mm, ft = 0.25 mm, Merck Germany). Different solvent mixtures were used (S1/hexane:dichloromethane (9:1), S2/hexane:ethyl acetate (8:2), S3/chloroform:methanol (8:2) and S4/chloroform:methanol (6:4)) and chromatograms were visualized by exposure under UV254/365 light, and by spraying with p-anisaldehyde and vanillin sulfuric reagents followed by heating [20, 21].

Fractionation of Ea extract and isolation of compounds

According to the results of the BSL screening assay (Table 1), ethyl acetate extract (Ea) was selected as the most active (LC = 249.26 μg mL⁻¹) compared to other extracts (Pe, Ch, Me) and, hence, was subjected to chromatographic fractionation. The Ea extract (6 g) was dissolved into small volume (5 ml) of ethyl acetate and the solution was subjected to a LC column (26 mm diameter × 300 mm height) filled with silica gel (type 60Å, 230-400 mesh ASTM, Merck). Elution was carried out with mixtures of hexane and ethyl acetate (90:10[®]70:30) of increasing polarity as described in Figure 1. Fifteen (50) fractions (100 ml each) were collected.

After TLC control using the different solvent systems S1 to S4 and staining with vanillin sulphuric and p-anisaldehyde reactifs [20], chromatographic fractions with similar TLC patterns were combined to afford sixteen (16) fractions

(F₁-F₁₆), as shown in Figure 1. Subsequent BSL assay screening revealed that the fractions F₄ (56.3 mg), F₈ (37.9 mg), F₉ (259.6 mg), F₁₁ (134.9 mg) and F₁₆ (862.4 mg) were highly active (LC<100 µg mL⁻¹).




Figure 1. Flow chart of BSL assay-guided fractionation of cytotoxic ethyl acetate extract of *Anacyclus pyrethrum* roots

A phytochemical screening showed a homogeneity in the composition of the fraction F₁₁ (LC = 42.5 µg mL⁻¹), with mainly alkalimides compounds as revealed upon staining with anisaldehyde reagent [20]. Fraction F₁₁ (LC = 42.5 µg mL⁻¹) was subsequently applied to a mini LC column chromatography (15 mm diameter × 200 mm height), filled with silica gel (high purity grade 40, 70-230 mesh ASTM, Merck), and eluted with mixtures of increasing polarity of cyclohexane and diethylether (100:0 to 100:100), as described in Figure 1. Thirty six (36) subfractions (F_{11.1} to F_{11.36}) were obtained and monitored by TLC. Alkalimides were purified by preparative TLC silica gel plates (20×20, Merck) using solvent system S2 from subfraction F_{11.7} (compounds II and III with 12.3 mg and 5.6 mg) and from Subfraction F_{11.9} (compounds I and IV with 7.7 mg and 3.8 mg). Compounds purity was assessed by TLC silica gel using three solvent mixtures of different polarity: hexane:ethyl acetate (2:1), Hexane: chloroform:formic acid (2:6:1) and petroleum ether:ethylacetate:formic acid (7:30:1).

Structural identification

The structures of compounds (I-VI) were characterized by 1H-NMR (BRÜKER Avance 300 MHz, Salerno University, Italy) and comparison with relevant data from bibliography. Samples were dissolved in CDCl₃.

Compound (I): White crystals; 1H-NMR δ (CDCl₃, 300 MHz) ppm: 0.86 (3H, m, H-10), 0.88 (6H, d, H-3'), 1.32 (6H, m, H-7, H-8 and H-9), 1.88 (1H, m, H-2'), 2.10 (2H, m, H-6), 3.20 (2H, m, H-1'), 5.00 (1H, br s, N-H), 5.77 (1H, m, H-2), 6.25 (1H, m, H-5); 6.75 (1H, m, H-4); 6.96 (1H, m, H-3).

Compound (II): White crystals; 1H-NMR δ (CDCl₃, 300 MHz) ppm: 0.92 (6H, t, H-3' and H-12), 1.26 (8H, m, H-2', H-6, H-5 and H-11), 1.54 (2H, dd, H-7), 1.85 (2H, m, H-10), 2.19 (2H, m, H-4), 3.17 (2H, d, H-1'), 5.38 (1H, br s, N-H), 5.79 (1H, m, H-2), 6.10 (1H, m, H-9); 6.73 (1H, m, H-8); 6.92 (1H, m, H-3).

Compound (III): White crystals; 1H-NMR δ (CDCl₃, 300 MHz) ppm: 0.91 (6H, d, H-3'), 0.94 (3H, m, H-14), 1.28 (14H, m, H-7, H-8, H-9, H-10, H-11, H-12 and H-13), 1.74 (1H, dd, H-2'), 2.36 (2H, t, H-6), 3.20 (2H, m, H-1'), 5.41 (1H, br s, N-H), 5.78 (1H, m, H-2), 6.18 (1H, m, H-5); 6.83 (1H, m, H-4); 6.98 (1H, m, H-3).

Compound (IV): White crystals; 1H-NMR δ (CDCl₃, 300 MHz) ppm: 0.94 (6H, m, H-3' and H-9), 1.30 (4H, d, H-2' and H-8), 1.85 (2H, m, H-7), 2.73 (2H, m, H-4), 3.21 (2H, d, H-1'), 5.02 (1H, br s, N-H), 5.80 (1H, m, H-2), 6.24 (1H, m, H-6); 6.83 (1H, m, H-5); 7.00 (1H, m, H-3).

Brine shrimp lethality (BSL) assay

Eggs of brine shrimp (*Artemia salina*) (Sera®, Bosnia-Herzegovina) were hatched in aqueous brine solution (3.8 g/L at 27-28°C). After 48 hours, the brine shrimp nauplii freed from egg shells were ready for the assay.

The BSL assay was conducted as described earlier with some modifications [22-25]. Briefly, pellitory samples (crude extracts and fractions) were tested at various concentrations (1-1000 µg/mL) in a brine solution containing 1% DMSO (v/v). A suspension solution of 10 nauplii were drawn through a glass capillary and placed in each compartment of a 12-well microplate. Negative control wells contained 1% DMSO in salty water. Pellitory solutions and control were tested in tetraplicate (n=4).

After 24h, the microplates were then examined under light, using a dissection microscope. The number of dead larvae was counted. Nauplii were considered dead if they did not exhibit any internal or external movement during several seconds of observation [26,27].

The percentage of mortality (% M) was calculated by the following formula as previously reported [28]:

$$(\% \text{ lethality}) = \frac{\text{Total nauplii} - \text{alive nauplii}}{\text{Total nauplii}} \times 100$$

Lethal concentration (LC₅₀) values were determined Finney's Probit regression analysis [29]. The lethality of the extracts on the brine shrimp was classified as previously reported [30,31]: LC₅₀ < 1000 µg/mL was "toxic", LC₅₀ = 500-1000 µg/mL was "weakly toxic", LC₅₀ = 100-500 µg/mL was "moderately toxic", and LC₅₀ < 100 µg/mL was "strongly toxic".

Statistics analysis


The data were collected from repeated measures (n = 4), and the results were reported as the mean ± standard deviation (SD). Data were analysed using Microsoft Excel 2007 (Redmond, WA, USA) and Microcal 6.0 (Microcal Software®, Inc.) for windows. The level of statistical significance between treated and untreated groups (control) was assessed by Tukey's test. The significance threshold was set at p < 0.05.

RESULTS

Pellitory roots extraction with solvents of different polarities yielded extractive materials of different physical consistencies. Both chloroform (Ch) and ethyl acetate (Ea) extracts were of powdery aspect, while those of methanol (Me) and petroleum ether (Pe) were greasy to resinous. Extracts are expressed with by yields, and are reported with their respective cytotoxic activity on the Brine Shrimp Lethality (BSL) assay in Table 1.

The Ae extract of the Pellitory treated group showed 100% mortality at the concentration of 500 $\mu\text{g/mL}$. The remaining extracts (Pe,Ch,Me) treated groups exhibited no or low mortality in the range of tested concentrations (1-1000 $\mu\text{g/mL}$). No lethality was found in the negative control (DMSO) group.

The BSL screening of Pellitory extracts showed that the most nauplii mortality was localized in the ethyl acetate extract (Ea), which displayed an $\text{LC}_{50} = 249.3 \mu\text{g/mL}$. To further investigate the active Ea extract, BSL assay-guided chromatographic fractionation (Figure 1) led to sixteen fractions, among which fraction F11 was revealed to be highly cytotoxic, with $\text{LC}_{50} = 42.5 \mu\text{g/mL}$. Multistep purifications of the active F11 afforded four compounds (I-IV) (Figure 2).



Compound names	R ₁	R ₂
(I) N-isobutyldeca-2,4-dienamide (Pellitorine)	$\text{CH}_3(\text{CH}_2)_4\text{-CH=CH-}$	$-\text{CH}(\text{CH}_3)_2$
(II) N-propyldodeca-2,8-dienamide	$\text{CH}_3(\text{CH}_2)_2\text{-CH=CH}(\text{CH}_2)_4\text{-}$	$-\text{CH}_2\text{CH}_3$
(III) N-isobutyltetradeca-2,4-dienamide	$\text{CH}_3(\text{CH}_2)_6\text{-CH=CH-}$	$-\text{CH}(\text{CH}_3)_2$
(IV) N-propylnona-2,5-dienamide	$\text{CH}_3(\text{CH}_2)_2\text{-CH=CH-CH}_2\text{-}$	$-\text{CH}_2\text{CH}_3$

Figure 2. Alkamides and their chemical structures

Compound I was isolated as white crystals. The 1-H NMR spectrum exhibited characteristic signals for a N-isobutylamide group at δ 3.20 ppm (2H,m, H-1'), δ 1.88 ppm (1H,m, H-2') and δ 0.88 ppm (6H,d, H-3'), together with amide proton signal at δ 5.00 ppm (1H, br s) as previously reported for isobutylamide groupe [32]. In the low field region four olefinic methines signals (1H, m) are observed at δ 5.77 (H-2), 6.25(H-5), 6.75(H-4) and 6.96 (H-3) ppm. Multiplet signals at δ 1.32 (6H) and 2.10 (2H) are compatible with the germinal protons (H-7,H-8 and H-9) and H6, respectively. High field signals at δ 0.86 (3H, t) is compatible with one methyl protons (H-10). The structure of compound I is assigned to the well-known pellitorine or N-isobutydeca-2,4-dienamide, on the basis of H-NMR data which were in full agreement with literature values [33-38].

Compound II was isolated as white crystals. The 1-H NMR spectrum exhibited signals for a N-propylamide group: δ 3.17 ppm (2H, H-1'), δ 1.26 ppm (2H, H-2') and δ 0.92 ppm (3H, H-3'), together with that at δ 5.38 ppm (1H, br s) assigned to the amide proton. Four olefinic methines signals were observed at δ 5.79, 6.10, 6.73 and 6.92 ppm (1H each) as previously reported for isolated

double bonds in alkamides [39]. On the basis of this reference, we deduced the presence of two double bonds in position 2 and 8. By comparing the 1H NMR data with previously published data, compound II was identified as: N-propyldodeca-2,8-dienamide.

Compound III was isolated as white crystals. The 1-H NMR spectrum exhibited characteristic signals for a N-isobutylamide group at δ 3.20 ppm (2H), δ 1.74 ppm (1H) and δ 0.91 ppm (6H), together with amide proton signal at δ 5.41 ppm (1H, br s) as previously reported [40]. Four olefinic methine signals are observed at δ 5.78, 6.18, 6.83 and 6.98 ppm (1H each), which are compatible with the presence of two double bonds conjugated with the amide carbonyl as previously recorded for alkamides [32,40]. On the basis of previously 1-H NMR published data, compound III is identified as: N-isobutyltetradeca-2,4-dienamide.

Compound IV was isolated as white crystals. The 1-H NMR spectrum was some points similar to that of compound II. It shows signals for a N-propylamide group: δ 3.21 ppm (2H), δ 1.30 ppm (2H) and δ 0.94 ppm (3H), together with that at δ 5.41 ppm (1H, br s) assigned to the amide proton as previously reported [40]. The four olefinic methine signals were observed at δ 5.80, 6.24, 6.83 and 7.00 ppm (1H), as recorded for alkamides with double conjugations in position 2 and 5 [40]. By comparing the 1-H NMR data with previously published data [32,40], compound 3 was identified as: N-propylnona-2,5-dienamide.

DISCUSSION

Pellitory (*Anacyclus pyrethrum*) roots has been known since ancient times and has been and is used as medicinal plant applied – among other ailments – against cancer [8]. Of special pharmacological interest, pellitory produces several N-alkamides [14], which are secondary metabolites in plants. Because of their wide structural diversity, these compounds have attracted several research groups to study their pharmacological behaviours [41]. Numerous reports have dealt with the anesthetic, analgesic and anti-inflammatory, but also with the anticholinesterase, antidiabetic, antiparasitic, anticancer, molluscicidal, antiprotozoal and insecticidal activities of N-alkamides [36,41,42].

Among the multiple pharmacological effects of alkamides, its anticancer property is the most promising. Thus, many alkamides isolated from the Asteraceae and other botanical families were shown to exhibit *in vitro* cytotoxic activity against cancer cell lines [7,41,42].

Worth noting is the pellitorine (occurring in pellitory roots) has been shown to have strong cytotoxic activities against HL60 (Human promyelocytic leukemia strain) and MCT-7 (Brest cancer strain) cell lines, with IC_{50} values of 13.0 $\mu\text{g/mL}$ and 1.8 $\mu\text{g/mL}$, respectively [43]. Piplartine and analogues have exhibited potent effects in human breast carcinoma MCF-7 cells, whilst being relatively non-toxic to non-tumorigenic MCF-10a cells [44,45]. Other alkamides, as capsaicin was reported to induce the apoptosis of prostate cancer cell lines [46], while pharnilatin did so for those of skin melanoma, as well as lung, ovary and colon [47]. These reports and many involving others alkamides, make these

type of compounds a relatively new and promising group of natural products as source of new anticancer agents [41].

BSL assay is recognized as a general test for bioactive compounds screening [19,48-53]. The technique is described as easily mastered, costs little, and utilizes small amount of test material [50]. Since its introduction in 1982, by Meyer and al. [30], it appears that BSL test is predictive of cytotoxicity and has been successively employed for bioassay-guided fractionation of active cytotoxic and antitumor agents as reported by several authors [48,52-56]. A significant correlation between the brine shrimp assay and *in vitro* growth inhibition of human solid tumor cell lines has been demonstrated by the National Cancer Institute (NCI, USA) [51].

In our toxicity evaluation of pellitory root's extracts (Pe, Ch, Ea, Me), the BSL test showed that the most of the nauplii mortality was localized in the ethyl acetate extract (Ea), which displayed an LC₅₀ = 249.3 µg/mL. This value is considered as "toxic" [30,31], and hence bioguided fractionation of Ea active extract of *Anacyclus pyrethrum* root was undertaken and revealed a highly cytotoxic fraction (F₁₁) with LC₅₀ of 42.5 µg/mL. Multistep purifications of the active (F₁₁) fraction have afforded four alkalimides (Figure 2): namely N-isobutyldeca-2,4-dienamide or pellitorine (I), N-propyldodeca-2,8-dienamide (II), N-isobutyltetradeca-2,4-dienamide (III) and N-propylnona-2,5-dienamide (IV). The compounds (I) and (III) were previously reported from *A. pyrethrum* roots and other species [14]. The two remaining alkalimides (II) and (IV) are isolated for the first time from pellitory, but were cited previously in the natural alkalimides database [41].

In our study, the BSL test appears to be effective in the detection of potentially toxic fractions containing N-alkalimides, such as pellitorine a well known cytotoxic agent [43]. Hence, our approach of using the *Artemia salina* mortality test to guide the fractionation of the pellitory root Ea extract was productive, and thus supports its appropriateness as a pre-screening tool for cytotoxic compounds.

CONCLUSION

Pellitory roots have been cited as a cancer remedy in folk medicine. Recent study have shown that pellitory alcoholic extract have cytotoxic activity against human colorectal cancer cell lines. Our study suggests that cytotoxic activity is localized in the ethyl acetate extract (Ea) of pellitory root, and this effect might be associated with the presence of alkalimides, pellitorine and its analogues (II-IV). Although pellitorine (I) has previously shown strong cytotoxic activities against human cancer cell lines, no studies have been reported on cytotoxic activities of other isolated compounds (II to IV). Subsequently, these alkalimides will be considered in future study as candidates for in depth *in vitro* evaluation of their cytotoxicity against cancer and normal cell lines. Finally, through this study, BSL assay demonstrate again its usefulness as bench-top assay in exploring plant extracts for cytotoxic compounds.

REFERENCES

1. Kishor KV, Lalitha GK. Pharmacognostical studies on the root of *Anacyclus pyrethrum* DC. *Indian Journal of Natural Products and Resources*. 2012;3:518-26.
2. Puri HS. *Rasayana – Ayurvedic herbs for longevity & rejuvenation*. Vol. 2. London: Taylor & Francis; 2003:20-1.
3. Sharma V, Thakur M, Chauhan NS, Dixit VK. Effects of petroleum ether extract of *Anacyclus pyrethrum* DC. on sexual behavior in male rats. *Journal of Chinese Integrative Medicine*. 2010;8(8):767-72.
4. Pahuja M, Mehla J, Reeta KH, Joshi S, Gupta YK. Root extract of *Anacyclus pyrethrum* ameliorates seizures, seizure-induced oxidative stress and cognitive impairment in experimental animals. *Epilepsy Research*. 2012;98(2-3):157-65.
5. Pahuja M, Mehla J, Reeta KH, Tripathi M, Gupta YK. Effect of *Anacyclus pyrethrum* on pentylentetrazole-induced kindling, spatial memory oxidative stress and rho-kinase II expression in mice. *Neurochem Res*. 2013;38(3):547-56.
6. Veyer L, Taevernier L, Roche N, Peremans K, Burvenich C, De Spiegeleer B. Quantitative transdermal behavior of pellitorine from *Anacyclus pyrethrum* extract. *Phytomedicine*. 2014;21(14): 1801-07.
7. Usmani A, Khushtar M, Arif M, Siddiqui MA, Sing SP, Mujahid M. Pharmacognostic and phytopharmacology study of *Anacyclus pyrethrum*: an insight. *Journal of Applied Pharmaceutical Science*. 2016;6(3):144-50.
8. Singh A. *Compendia Of World's Medicinal Flora*. Boca Raton: CRC Press; 2006: 58-9.
9. Bellakhdar J. *La pharmacopée marocaine traditionnelle: Médecine arabe ancienne et savoirs populaires*. Paris: Ibis Press; 1997.
10. Boulos L. *Medicinal plants of North Africa*. Michigan: Reference Publications, Inc; 1983.
11. IUCN. *A guide to medicinal plants in North Africa*. Malaga: Centre for Mediterranean Cooperation; 2005.
12. Sujiet K, Darwin CR, Suba V. Antioxidant activity of ethanolic root extract of *Anacyclus pyrethrum*. *International Research Journal of Pharmacy*. 2011;2(12):222-6.
13. Singh DK, Nirwan S, Babbar SB. Micropropagation of *Anacyclus pyrethrum* and chemical profiling of the regenerated plants for pellitorine, the active principle. *Plant Cell Tiss Organ Cult*. 2015; 122(1):249-55.
14. Boonen J, Sharma V, Dixit VK, Burvenich C, De Spiegeleer B. LC-MS N-alkylamide profiling of an etanolic *Anacyclus pyrethrum* root extract. *Planta Med*. 2012;78:1784-95.
15. Hinz B, Woelkart K, Bauer R. Alkalimides from *Echinacea* inhibit cyclooxygenase-2 activity in human neuroglioma cells. *Biochemical and Biophysical Research Communications*. 2007;360(2): 441-6.
16. Shahraki MR, Shahraki S, Arab MR, Shahrakipour M. The Effects of Aqueous Extract of *Anacyclus pyrethrum* on sperm count and reproductive organs in adult male rats. *Zahedan Journal of Research in Medical Sciences*. 2015;17(2):42-6.
17. Mohammadi A, Mansoori B, Chaghakaboodi K, Baradaran B. Cytotoxic effects of *Anacyclus pyrethrum* plant extract in oral cancer cell (KB cell line). *J Urmia Univ Med Sci*. 2016;27(4): 257-65.
18. Mohammadi A, Mansoori B, Baradaran PC, Baradaran SC, Baradaran B. *Anacyclus Pyrethrum* extract exerts anticancer activities on the human colorectal cancer cell line (HCT) by targeting apoptosis, metastasis and cell cycle arrest. *J Gastrointest Cancer*. 2017;48(4):333-40.
19. McLaughlin JL. *Assays for bioactivity*. In: Hostettmann K (Ed), *Methods in Plant Biochemistry*, Vol.6, London: Academic Press; 1991:1-33.
20. Wagner H, Bladt S. *Plant drug analysis: a thin layer chromatography atlas*. 2 edition. Berlin: Springer-Verlag; 2011.
21. Harborne JB. *Phytochemical methods. A guide to modern techniques of plant analysis*. London: Chapman and Hall Ltd; 1984.
22. Bidau CJ, Amat AG, Yajia ME, Marti DA, Gimenez MD, Riglos AG and al. Evaluation of cytotoxic and mitodepressive activity of aqueous extracts from thirteen Argentine medicinal plants. *Acta Farm. Bonaerense*. 2006;25(4):555-9.

23. Lagnika L, Anago E, Sanni A. Screening for antibacterial, antioxidant activity and toxicity of some medicinal plants used in Benin folkloric medicine. *Journal of Medicinal Plants Research*. 2011;5(5):773-7.
24. Syahmi AR, Vijayarathna S, Sasidharan S, Latha LY, Kwan YP, Lau YL and al. Acute oral toxicity and brine shrimp lethality of *Elaeis guineensis* Jacq., (oil palm leaf) methanol extract. *Molecules*. 2010; 15(11):8111-21.
25. Salawu KM, Ajaiyeoba EO, Ogbale OO, Adeniji JA, Faleye TC, Agunu A. Antioxidant, brine shrimp lethality and antiproliferative properties of gel and leaf extracts of *Aloe schweinfurthii* and *Aloe vera*. *Journal of Herbs, Spices & Medicinal Plants*. 2017;23(4):1-10.
26. Pisutthanana S, Plianbangchang P, Pisutthanana N, Ruanruaya S, Muanrita O. Brine shrimp lethality activity of Thai medicinal plants in the family Meliaceae. *Naresuan University Journal*. 2004; 12(2):13-8.
27. Sahgal G, Ramanathan S, Sasidharan S, Mordi MN, Ismail S, Mansor SM. Brine shrimp lethality and acute oral toxicity studies on *Swietenia mahagoni* (Linn.) Jacq. seed methanolic extract. *Pharmacognosy Research*. 2010;2(4):215-20.
28. Hartl M, Humpf HU. Toxicity assessment of fumonisins using the brine shrimp (*Artemia salina*) bioassay. *Food and Chemical Toxicology*. 2000;38(12):1097-102.
29. Hamidi MR, Jovanova B, Panovska TK. Toxicological evaluation of the plant products using brine shrimp (*Artemia salina* L.) model. *Macedonian Pharmaceutical Bulletin*. 2014;60(1):9-18.
30. Meyer BN, Ferrigni NR, Putnam JE, Jacobsen LB, Nichols DE, McLaughlin JL. Brine shrimp: a convenient general bioassay for active plant constituents. *Planta Medica*. 1982;45(5):31-4.
31. Oladimeji HO, Nia R, Essien EE. *In vitro* antimicrobial and brine shrimp lethality potential of the leaves and stem of *Calotropis procera* (Ait). *African Journal of Biomedical Research*. 2006;9:205-11.
32. Rosario SL, Da Silva AJ, Parente JP. Alkamides from *Cissampelos glaberrima*. *Planta Med*. 1996;62(4):376-7.
33. Rios-Chavez P, Ramirez-Chavez E, Armenta-Salinas C, Torres-Molina J. *Acmella radicans* var. *radicans*: *In vitro* culture establishment and alkamide content. *In vitro Cellular & Developmental Biology-Plant*. 2003;39:37-41.
34. Wu S, Sun C, Pei S, Lu Y, Pan Y. Preparative isolation and purification of amides from the fruits of *Piper Longum* L. by upright counter – current chromatography and reversed – phase liquid chromatography. *Journal of Chromatography*. 2004;1040(2):193-204.
35. Olah Z, Redei D, Pecze L, Vizler C, Josvay K, Forgo P and al. Pellitorine, an extract of *Tetradium daniellii*, is an antagonist of the ion channel TRPV1. *Phytomedicine*. 2017;34: 44-9.
36. Althaus JB, Kaiser M, Brun R, Schmidt TJ. Antiprotozoal activity of *Achillea ptarmica* (Asteraceae) and its main alkamide constituents. *Molecules*. 2014;19(5): 6428-38.
37. Cheng YB, Liu RH, Ho MC, Wu TY, Chen CY, Lo IW and al. Alkamides of *Acmella oleracea*. *Molecules*. 2015;20:6970-7.
38. Silveira N, Saar J, Santos AD, Barison A, Sandjo LP, Kaiser M and al. A new alkamide with an Endoperoxide structure from *Acmella ciliate* (Asteraceae) and its *in vitro* antiplasmodial activity. *Molecules*. 2016;21(6):765-70.
39. Elazzouzi H, Khennouchi S, Bentayeb A, Elhilali F, Zair T. Effets biocides des alcaloides extraits des racines d'*Anacyclus pyrethrum* L. (Asteracées) sur *Callosobruchus maculatus* (Fab.) (Coleoptera: Bruchidae). *International Journal of Innovation and Applied Studies*. 2015;13(4):756-74.
40. Chen Y, Fu T, Tao T, Yang J, Chang Y, Wang M and al. Macrophage activating effects of new alkamides from the roots of *Echinacea* species. *J. Nat. Prod*. 2005;68(5):773-6.
41. Boonen J, Bronselaer A, Nielandt J, Veryser L, De Tré G, De Spiegeleer B. Alkamid database: Chemistry, occurrence and functionality of plant N-alkylamides. *J Ethnopharmacol*. 2012;142 (3):563-90.
42. Rios MY. Natural Alkamides: Pharmacology, Chemistry and Distribution. In: *Drug Discovery Research in Pharmacognosy*, Omboon Vallisuta and Suleiman M. Olimat (eds). China: InTech; 2012:107-44.
43. Ee GC, Lim CM, Rahmani M, Shaari K, Bong CF. Pellitorine, a potential anti-cancer lead compound against HL60 and MCT-7 cell lines and microbial transformation of piperine from *Piper Nigrum*. *Molecules*. 2010;15(4):2398-404.
44. Bezerra DP, Pessoa C, de Moraes MO, Saker-Neto N, Silveira ER, Costa-Lotuf LV. Overview of the therapeutic potential of pipartine (piperlongumine). *European Journal of Pharmaceutical Sciences*. 2013;48(3):453-63.
45. Meegan MJ, Nathwani S, Twamley B, Zisterer DM, O'Boyle NM. Piperlongumine (Piplartine) and Analogues: Antiproliferative Microtubule-Destabilising Agents. *European Journal of Medicinal Chemistry*. 2017;125:453-63.
46. Luo XJ, Peng J, Li YJ. Recent advances in the study on capsaicinoids and capsinoids. *European Journal of Pharmacology* 2011;650:1-7.
47. Kim KH, Choi SU, Son MW, Lee KR. Two new phenolic amides from the seeds of *pharbitis nil*. *Chemical and Pharmaceutical Bulletin*. 2010;58:1532-5.
48. Carballo JL, Hernández-Inda ZL, Pérez P, García-Grávalos MD. A comparison between two brine shrimp assays to detect *in vitro* cytotoxicity in marine natural products. *BMC Biotechnol*. 2002;2:17-21.
49. Harwig J, Scott PM. Brine shrimp (*Artemia salina* L.) larvae as a screening system for fungal toxins. *Appl Microbiol*. 1971;21(6): 1011-6.
50. McLaughlin JL, Rogers LL. The use of biological assays to evaluate botanicals. *Drug Information Journal*. 1988;32:513-24.
51. Anderson JE, Goetz CM., McLaughlin JL, Suffness M. A blind comparison of simple benchtop bioassay and human tumour cell cytotoxicities as antitumor prescreens. *Phytochem Analysis*. 1991;2:107-11.
52. Solís PN, Wright CW, Anderson MM, Gupta MP, Phillipson JD. A microwell cytotoxicity assay using *Artemia salina*. *Plant Med*. 1993;59:250-2.
53. Mackeen MM, Ali AM, Lajis NH, Kawazu K, Hassan Z, Amran M and al. Antimicrobial, antioxidant, antitumour-promoting and cytotoxic activities of different plant part extracts of *Garcinia atroviridis* Griff. Ex T. Anders. *J Ethnopharmacol*. 2000;72:395-402.
54. Zhao G, Hui Y, Rupprecht JK, McLaughlin JL, Wood KV. Additional bioactive compounds and trilobacin, a novel highly cytotoxic acetogenin, from the bark of *Asimina triloba*. *Journal of Natural Products*. 1992;55:347-56.
55. Rieser MJ, Gu ZM, Fang XP, Zeng L, Wood KV, McLaughlin JL. Five novel mono-tetrahydrofuran ring acetogenins from the seeds of *Annona muricata*. *Journal of Natural Products*. 1996;59: 100-8.
56. Mongelli E, Pomilio AB, Sanchez JB, Guerra FM, Massanet GM. ent-Kaur-16-en-19-oic acid, a KB cells cytotoxic diterpenoid from *Elaeoselinum foetidum*. *Phytotherapy Research*. 2002;16:387-8